
All patients taking antipsychotics are at risk for Tardive Dyskinesia

NO MATTER THE ANTIPSYCHOTIC, NO MATTER THE INDICATION¹⁻³



FGA, first-generation antipsychotic;

SGA, second-generation antipsychotic.

N=11,493, meta-analysis including 41 studies that provided cross-sectional data from at least 15 FGA-, SGA-, or FGA + SGA-treated subjects and used a standardized rating scale to evaluate TD **Bipolar disorder**


Schizophrenia

Major depressive disorder

Anxiety disorder

LEARN HOW TO TELL THE DIFFERENCE

Tardive Dyskinesia



"Look for more than just the presence of a movement the type of movement is just as important for diagnosis."

-Desiree M. Matthews, PMHNP-BC Nurse Practitioner Outpatient Behavioral Health Advanced Practice Provider Clinical Liaison, Monarch Charlotte, NC Paid consultant of Neurocrine Biosciences, Inc.

References

1. Carbon M, et al. J Clin Psychiatry. 2017;78(3):e264-e278. 2. Citrome L, et al. Clin Ther. 2013;35(12):1867-1875. 3. Leslie DL, et al. Psychiatr Serv. 2009;60(9):1175-1181. 4. McEvoy J, et al. Qual Life Res. 2019;28(12):3303-3312. 5. Ward KM, Citrome L. Neurol Ther. 2018;7(2):233-248. 6. Savitt D, Jankovic J. J Neurol Sci. 2018;389:35-42. 7. Caroff SN, et al. J Clin Psychiatry. 2020;81(2):19cs12983. 8. Tarsy D. Curr Treat Options Neurol. 2000;2(3):205-214. 9. Guy W. ECDEU Assessment Manual for Psychopharmacology. US Department of Health, Education, and Welfare, Public Health Service, Alcohol, Drug Abuse, and Mental Health Administration, National Institute of Mental Health, Psychopharmacology Research Branch, Division of Extramural Research Programs; 1976. 10. American Psychiatric Association. Diagnostic and Statistical Manual of Mental Disorders. 5th ed. American Psychiatric Association; 2013. 11. Mathews M, et al. Psychiatry. 2005;2(3):36-41. This content was sponsored and co-developed by Neurocrine Biosciences, Inc.



MIND[•]TD[•] MED-MSL-TD-US-0248 v2 CP-TD-US-0664 v3 09/2022